

Original Research

Meaning in Life: A Novel Factor for Promoting Well-being in COPD

Caitlin Batzlaff, MD^{1,2} Madison Roy, MS² Johanna Hoult, MA² Roberto Benzo, MD^{1,2}

Abstract

Introduction: Research evidence indicates that meaning in life (MIL) is a meaningful outcome in individuals living with chronic illness, but evidence is lacking for chronic obstructive pulmonary disease (COPD).

Objective: We hypothesized that MIL is independently associated with clinically meaningful outcomes for patients with COPD.

Methods: We performed cross-sectional analysis from a large cohort with moderate–severe COPD that participated in a home pulmonary rehabilitation study. MIL was measured using the Meaning in Life Questionnaire (MLQ), with domains of Presence and Search. The study included 340 participants: mean age 69 years old (standard deviation [SD] 9.4), 55.3% female, forced expiratory volume in 1 second (FEV₁) 44.3% (SD 19.4), and a modified Medical Research Council (mMRC) dyspnea scale of 2.69 (SD 0.84).

Results: Patients with high MLQ Presence (versus low score) had meaningfully better scores (beyond the minimal clinically important difference) across all Chronic Respiratory Questionnaire (CRQ) domains: Dyspnea, Fatigue, Emotions, and Mastery ($p \leq 0.02$); self-management ($p \leq 0.001$); social support ($p \leq 0.001$); anxiety ($p \leq 0.001$); and depression ($p \leq 0.01$) scores. When adjusting for age, sex, FEV₁, mMRC, social support, and anxiety, MLQ Presence was independently associated with CRQ domains Fatigue, Emotions, and self-management ($p \leq 0.01$). MLQ Search was independently associated with CRQ Dyspnea, Fatigue, and Mastery ($p < 0.05$).

Conclusion: In patients with COPD, the perception that life has meaning or the willingness to search for MIL is associated with the outcomes that patients consider most important. Our results are novel as MIL is a potentially modifiable outcome that could complement person-centered conversations during clinical visits, pulmonary rehabilitation, and health coaching.

1. Department of Pulmonary and Critical Care Medicine, Mayo Clinic, Rochester, Minnesota, United States

2. Mindful Breathing Lab, Mayo Clinic, Rochester, Minnesota, United States

Abbreviations:

CI=confidence interval; **COPD**=chronic obstructive pulmonary disease; **CRQ**=Chronic Respiratory Questionnaire; **FEV₁**=forced expiratory volume in 1 second; **GAD-2**=Generalized Anxiety Disorder 2-item; **GOLD**=Global initiative for chronic Obstructive Lung Disease; **ISEL**=Interpersonal Support Evaluation List-12; **MCID**=minimal clinically important difference; **MIL**=meaning in life; **MLQ**=Meaning in Life Questionnaire; **MLQ-P**=presence of meaning; **MLQ-S**=search for meaning; **mMRC**=modified Medical Research Council; **PHQ-9**=Patient Health Questionnaire; **SD**=standard deviation; **SMAS-30**=Self-Management Ability Scale

Funding Support:

National Heart, Lung, and Blood Institute (K24 HL138150).

Citation:

Batzlaff C, Roy M, Hoult J, Benzo R. Meaning in life: a novel factor for promoting well-being in COPD. *Chronic ObstrPulm Dis*. 2024; 11(4):341-349. doi: <https://doi.org/10.15326/jcopdf.2023.0476>

Publication Dates:

Date of Acceptance: April 1, 2024

Published Online Date: May 6, 2024

Address correspondence to:

Roberto Benzo, MD
Mindful Breathing Laboratory
Division of Pulmonary, Critical Care, and Sleep Medicine
200 First Street SW
Rochester, MN 55905
Phone: (507) 293-1989
Email: Benzo.Roberto@mayo.edu

Keywords:

advocacy; care coordination; health-related quality of life; pulmonary rehabilitation; chronic airways obstruction

This article has an online supplement.

A portion of this work was presented as an oral presentation at the CHEST 2023 Annual Meeting in Honolulu, Hawaii.

Introduction

Meaning in life (MIL) is defined as “the sense made of, and significance felt regarding, the nature of one’s being and existence.”¹

MIL is associated with well-being in healthy populations¹ and there is some recent evidence that those who find meaning in life may be better able to cope with medical challenges.²⁻⁵ Justifiably, those attempting to process health-related challenges may prioritize defining life’s meaning to better cope.^{6,7} Prior studies have demonstrated that serious illness confronts one’s perception of life balance – their values, priorities, and future life endeavors. Adaptive thought processes invite complex and uncomfortable questions regarding meaning in life; allowing responses to be flexible and goals to be modified.⁸

MIL has been studied in various patient populations with chronic conditions, like cancer,⁹⁻¹² cardiac disease,¹³⁻¹⁶ and other life-threatening, chronic illnesses.^{3,5} MIL was related to higher psychological well-being in a sample of individuals living with spinal cord injury.¹⁷ Similarly, cancer patients with higher meaning in life reported improved quality of life,¹⁸ higher well-being,¹⁹ and lower levels of depressive symptoms and fatigue.²⁰

However, there is a knowledge gap in other prevalent conditions like chronic obstructive pulmonary disease (COPD). COPD is a leading cause of disability²¹ and has been in the top 10 since 1990. Living with COPD negatively impacts quality of life²²⁻²⁵ and further challenges MIL.²⁶

Given the growing importance of personalized support to patients to promote wellness while living with COPD, the inquiry of meaning in life may be significant during patient encounters to foster motivation for better coping.

We aimed to investigate the association of MIL with clinically meaningful outcomes for patients living with COPD in a large and well-characterized cohort.

Our study hypothesis was that the perception of MIL is independently associated with clinically meaningful outcomes for those living with COPD. Deepening our understanding of the association of MIL with outcomes may provide an opportunity to inform the conversations with patients about what is meaningful and worth pursuing, as well as setting goals for lifestyle change in chronic care interventions, like pulmonary rehabilitation, self-management programs, or health coaching.

Methods

Ethical Approval

This was a retrospective analysis using a large database from a randomized-controlled trial that received Mayo Clinic Institutional Review Board approval on January 11, 2018, IRB#17-009449.

Study Design and Setting

We conducted a cross-sectional analysis using baseline measures (before intervention) from a large cohort with moderate–severe COPD that participated in a home-based pulmonary rehabilitation study. Sites included: Mayo Clinic, Rochester, Minnesota; and Jacksonville, Florida; and Health Partners, Minneapolis–Saint Paul, Minnesota.

Inclusion Criteria

Inclusion criteria included a clinical diagnosis of COPD (primary inclusion criteria) confirmed by records, age ≥ 40 years old, a history of a minimum of 10 pack years of smoking, and the ability to communicate in English.

Participants

This analysis included 340 participants. Demographics data is displayed in Table 1.

Outcomes

Demographics information, forced expiratory volume in 1 second (FEV₁), and Global initiative for chronic Obstructive Lung Disease (GOLD)²⁷ group were gathered via chart review. To further assess their COPD, health-related quality of life metrics, and activity levels, the following information was collected: modified Medical Research Council (mMRC) dyspnea scale, Interpersonal Support Evaluation List-12 (ISEL), Patient Health Questionnaire (PHQ-9), Generalized Anxiety Disorder 2-item (GAD-2), Chronic Respiratory Questionnaire (CRQ), Meaning in Life Questionnaire (MLQ), and Self-Management Ability Scale (SMAS-30). Details can be found in the online supplement.

The MLQ is a 10-item tool with 2 separate subscales: the Presence of Meaning (MLQ-P) and the Search for Meaning (MLQ-S). Each subscale captures a different component of human functioning: presence as a tangible outcome versus search as an active and evolving process.¹

MLQ-P subscale represents the subjective sense that one’s life is meaningful. It refers to an individual’s understanding of themselves and how they fit into their surrounding environment or community; example “my life has a clear sense of purpose.”

MLQ-S subscale measures the drive, motivation, and intensity towards finding life’s purpose; example “I am looking for something that makes my life feel meaningful.”

Each subscale is measured by 5 items, rated using a 7-point Likert scale, with scores ranging from 5 to 35. Higher scores (≥ 24) are indicative of higher search or presence of meaning. Subscales had Cronbach’s alpha values 0.82–0.87 with one-month test-retest reliability of 0.70 (MLQ-P) and 0.73 (MLQ-S).¹

Table 1. Patient Demographics

Characteristic	Overall (N=340)
Age	
Mean (SD)	69.7 (9.4)
Sex	
Female	188 (55.3%)
Male	152 (44.7%)
mMRC	
Mean (SD)	2.7 (0.8)
Median	3.0
FEV₁	
Mean (SD)	44.3 (19.4)
GOLD Groups	
GOLD 1	15 (4.6%)
GOLD 2	101 (30.9%)
GOLD 3	127 (38.8%)
GOLD 4	84 (25.7%)
Race	
American Indian/Alaska Native	2 (0.6%)
Asian	2 (0.6%)
Black or African American	7 (2.1%)
More Than One Race	4 (1.2%)
Native Hawaiian or Other Pacific Islander	1 (0.3%)
White	324 (95.3%)
Ethnicity	
Hispanic or Latino	2 (0.6%)
Not Hispanic or Latino	328 (96.5%)
Unknown/Not Reported	10 (2.9%)
Population Density	
Urban	225 (66.6%)
Rural	113 (33.4%)
Marital Status	
Married	199 (58.5%)
Other	141 (41.5%)
Education	
High school or less	110 (32.5%)
Some college or more	228 (67.5%)

SD=standard deviation; mMRC=modified Medical Research Council; FEV₁=forced expiratory volume in 1 second; GOLD=Global initiative for chronic Obstructive Lung Disease

Data Analysis

Descriptive statistics were used to summarize patient demographics. Continuous data are presented as mean (standard deviation) and categorical data are presented as counts (percentages). Comparisons in quality-of-life measures between patients with high versus low MLQ-P and high versus low MLQ-S were made using Wilcoxon signed rank tests. Linear models were generated to determine the association between MLQ-P and MLQ-S with each of the CRQ domains and SMAS-30 total score. These models were adjusted for age, sex, FEV₁, ISEL, mMRC, GAD-2 and PHQ-9. Additional models were generated to investigate the association between MLQ-P and MLQ-S with PHQ-9. These models were adjusted for age, sex, FEV₁, ISEL, mMRC and GAD-2. Bootstrapping with 1000 replications

was conducted on all models to generate robust confidence intervals.^{28,29} *P* values <0.05 were considered statistically significant; analysis was completed using R.³⁰

Results

The study population was primarily White, non-Hispanic individuals with the majority of the participants having attended higher education. There was also a slight female predominance and 33% of the population came from rural areas. This demographic data is important to consider in the context of the below results, as they represent specific populations.

Individuals with high MLQ-P (versus low), have significantly better scores (beyond the minimal clinically

For personal use only. Permission required for all other uses.

important difference) across all CRQ domains: Dyspnea, Fatigue, Emotions, and Mastery ($p \leq 0.02$); self-management SMAS-30 ($p \leq 0.001$); social support ISEL ($p \leq 0.001$); anxiety GAD-2 ($p \leq 0.001$); and depression PHQ-9 ($p \leq 0.01$) scores (Figures 1-4).

As seen in Table 2 below, MLQ-P was independently associated with higher CRQ scores in the domains of Fatigue ($p < 0.001$) and Emotions ($p = 0.003$), as well as higher total self-management ability scores ($p \leq 0.001$).

MLQ-S was independently associated with higher CRQ scores in the domains of Dyspnea ($p = 0.007$), Fatigue ($p = 0.028$), and Mastery ($p = 0.041$) (Table 3).

Discussion

This study identified significant and independent associations of presence of and search for meaning in life to disease-specific quality of life, self-management abilities, and symptoms of depression in a well-characterized COPD cohort. Our results are novel as there is a scarcity of reports on the value of MIL in COPD. In the literature review, there was only one study identified that investigated meaning in life and self-care agency in the COPD population.³¹

High Presence of Meaning in Life

We found that those with high presence of MIL perceived better quality of life, better self-management abilities, better social support, and less anxiety and depression. The models adjusting for confounders show that presence (MLQ-P) was independently associated with higher scores in the CRQ domains of Fatigue, Emotions, Mastery, and total self-

management ability scores.

Our study confirmed previous findings that meaning in life predicts positive well-being in patients with chronic and/or life-threatening disease.^{3,7,9,32} Hsieh et al found that presence of meaning in life was a main mediator of depressive symptoms and trait mindfulness.⁹ Sherman et al identified that high personal meaning is independently associated with lower distress and improved coping in the primary care and cancer populations.⁷ Health anxiety is also reduced in those with high presence of meaning.² We could suggest that high presence of meaning in life may provide individuals with emotional resiliency and allows for better coping when faced with adversity.


On review of the cardiac literature, the capacity to find meaning in life after a significant cardiovascular event is associated with lower stress and improved mental health.¹³ Vos et al suggests a variety of mechanisms to elicit change and focus on the meaning-making,¹³ including adjustments to life perspective, connecting to fundamental values, building self-worth through hope and optimism, learning to self-regulate, and accepting limitations.

High Search for Meaning in Life

We found that the search for meaning in life (MLQ-S) was an independent predictor of higher CRQ scores in Dyspnea, Fatigue, and Mastery domains.

While a clear relationship between a high presence of meaning and healthy psychological well-being exists, the effect of the search for meaning is less established in the literature. Our results suggest the importance of searching for the meaning in life as it relates to the perception of well-

Figure 1. Comparing Those With Low Presence (Blue) to Those With High Presence (Orange) of Meaning in Life in Relation to Chronic Respiratory Questionnaire




*Denotes a p -value of 0.015

±Denotes a p -value of < 0.001

CRQ=Chronic Respiratory Questionnaire

For personal use only. Permission required for all other uses.


Figure 2. Comparing Those With Low Presence (Blue) to Those With High Presence (Orange) of Meaning in Life in Relation to the Self-Management Abilities Scale



±Denotes a p-value of <0.001

SMAS-30=Self-Management Abilities Scale

Figure 3. Comparing Those With Low Presence (Blue) to Those With High Presence (Orange) to Meaning in Life in Relation to the Generalized Anxiety Disorder 2-Item and Patient Health Questionnaire



*Denotes a p-value of <0.01

±Denotes a p-value of <0.001

GAD-2=Generalized Anxiety Disorder 2-item; PHQ-9=Patient Health Questionnaire

being in critical aspects of quality of life in people living with COPD: breathlessness (most common symptom), fatigue (second most common symptom), and self-management. The value of the search of meaning represents a novel discovery of this report that requires further validation. Dezutter et al distinguishes between adaptive and maladaptive search for MIL, suggesting an adaptive search outlook leads to more acceptance of chronic conditions.³ Our data supports


this concept in that those with a high search for meaning perceive less dyspnea and fatigue, while also having higher mastery scores, all critical outcomes in COPD.

Practical Implications

Addressing presence and search for meaning during clinical encounters in clinical practice and self-management programs (pulmonary rehabilitation and health coaching) would offer a complementary and personalized approach to

For personal use only. Permission required for all other uses.

Figure 4. Comparing Those With Low Presence (Blue) to Those With High Presence (Orange) to Meaning in Life in Relation to the Interpersonal Support Evaluation List-12



±Denotes a p-value of <math>< 0.001</math>

ISEL=Interpersonal Support Evaluation List-12

Table 2. Results of Linear Models^a and Bootstrapping Assessing the Relationship Between Meaning in Life Questionnaire Search and Primary Outcomes

Outcome	Effect of Having High Presence at Baseline ^b	P value	Adjusted R Square	95% Bootstrap CI
CRQ Dyspnea	-0.002	0.988	0.439	(-0.257, 0.274)
CRQ Fatigue	0.353	<math>< 0.001</math>	0.454	(0.138, 0.569)
CRQ Mastery	0.255	0.057	0.439	(-0.016, 0.503)
CRQ Emotions	0.269	0.003	0.643	(0.068, 0.445)
SMAS-30 Total	6.626	<math>< 0.001</math>	0.459	(4.399, 8.650)
PHQ-9 Score	-0.643	0.127	0.573	(-1.318, 0.069)

^aAdjusted for age, sex, FEV₁, ISEL, mMRC, GAD-2, and PHQ-9 (unless it is the outcome)

^bReference is low presence

CI=confidence interval; CRQ=Chronic Respiratory Questionnaire; SMAS-30=Self-Management Ability Scale; PHQ-9=Patient Health Questionnaire; FEV₁=forced expiratory volume in 1 second; ISEL=Interpersonal Support Evaluation List-12; mMRC=modified Medical Research Council; GAD-2=Generalized Anxiety Disorder 2-item

Table 3. Results of Linear Models^a and Bootstrapping Assessing the Relationship Between Meaning in Life Questionnaire Presence and Primary Outcomes

Outcome	Effect of Having High Search at Baseline ^b	P value	Adjusted R Square	95% Bootstrap CI
CRQ Dyspnea	0.376	0.007	0.400	(0.117, 0.643)
CRQ Fatigue	0.254	0.028	0.445	(0.044, 0.457)
CRQ Mastery	0.286	0.041	0.440	(0.049, 0.546)
CRQ Emotions	0.010	0.917	0.633	(-0.180, 0.207)
SMAS-30 Total	0.014	0.991	0.398	(-2.590, 2.521)
PHQ-9 Score	0.707	0.108	0.573	(-0.102, 1.688)

^aAdjusted for age, sex, FEV₁, ISEL, mMRC, GAD-2, and PHQ-9 (unless it is the outcome)

^bReference is low search

CI=confidence interval; CRQ=Chronic Respiratory Questionnaire; SMAS-30=Self-Management Ability Scale; PHQ-9=Patient Health Questionnaire; FEV₁=forced expiratory volume in 1 second; ISEL=Interpersonal Support Evaluation List-12; mMRC=modified Medical Research Council; GAD-2=Generalized Anxiety Disorder 2-item

For personal use only. Permission required for all other uses.

promote better coping, motivation, and wellness.³³

Our results provide novel evidence on the importance of meaning in life to sustain and even promote well-being in patients living with COPD. Our findings suggest that fostering high presence of meaning could lead to less fatigue, improved coping ability, and enhanced self-efficacy. While the search for meaning is associated with better dyspnea, fatigue, and sense of mastery.

Use of meaning-centered interventions has demonstrated positive outcomes in chronic diseases,^{10,19,34-38} but the literature is lacking in those with chronic obstructive pulmonary disease.

Our models for evaluating the associations of MIL with outcomes are robust (confirmed by bootstrapping) and a good explanation of the variance, as expressed for a high R-squared for a social science investigation. The addition of social support to the model aimed to respond to the concern from previous research regarding the interaction between MIL and social support. Liu et al discovered that those who had strong social support and improved self-management, as demonstrated by a decrease in care support needs, developed a stronger presence of meaning in life and quality of life post hospitalization.¹⁵

Limitations

Our study has limitations. First, despite multiple sites participating, the population was White, which may not be generalizable to other racial and/or ethnic groups. Second the cross-sectional analysis does not allow us to assess temporal or causal relations between the variables. Studies that investigate these relationships in a longitudinal fashion would allow for a better understanding of these relationships.

Conclusions

Our results innovatively suggest the impact of MIL on critical outcomes in individuals living with COPD. The dissemination of this work is important to increase awareness of the relationship between MIL and improved quality of life for those with COPD. It will stimulate research about MIL with the ultimate aim of promoting conversations about MIL during routine clinical visits, pulmonary rehabilitation and health coaching encounters, and self-management programs.

Acknowledgments

Author contributions: CB, MS, and RB contributed to the conception and/or design of the study, data analysis and/or interpretation, writing the article, and revisions prior to submission. JH contributed to the acquisition of data. All authors approved the final version of the manuscript submitted for publication. The authors would like to acknowledge the other members of the Mindful Breathing Lab: Maria Benzo, Josh Taylor, and Will Midthun for their contributions.

Data sharing statement: Individual participant data that underlie the results reported in this article, after deidentification, will be available. Study protocol, statistical analysis plan, and analytic code will not be available. Data will be available beginning 3 months and ending 48 months following article publication. Data will be shared with researchers who provide a methodologically sound proposal to achieve the aims of the approved proposal. Proposals should be directed to Benzo. Roberto@mayo.edu. To gain access, data requestors will need to sign a data access agreement.

Declaration of Interest

CB, MS, JH, and RB have no associations or conflicts of interest to disclose.

References

1. Steger MF, Frazier P, Oishi S, Kaler M. The meaning in life questionnaire: assessing the presence of and search for meaning in life. *J Couns Psychol.* 2006;53(1):80-93. <https://doi.org/10.1037/0022-0167.53.1.80>

2. Yek MH, Olendzki N, Kekecs Z, Patterson V, Elkins G. Presence of meaning in life and search for meaning in life and relationship to health anxiety. *Psychol Rep.* 2017;120(3):383-390. <https://doi.org/10.1177/0033294117697084>

3. Dezutter J, Casalin S, Wachholtz A, Luyckx K, Hekking J, Vandewiele W. Meaning in life: an important factor for the psychological well-being of chronically ill patients? *Rehabil Psychol.* 2013;58(4):334-341. <https://doi.org/10.1037/a0034393>

4. Krok D, Telka E, Zarzycka B. Illness perception and affective symptoms in gastrointestinal cancer patients: a moderated mediation analysis of meaning in life and coping. *Psychooncology.* 2019;28(8):1728-1734. <https://doi.org/10.1002/pon.5157>

5. Naghiyae M, Bahmani B, Asgari A. The psychometric properties of the Meaning in Life Questionnaire (MLQ) in patients with life-threatening illnesses. *Scientific World Journal.* 2020;2020:8361602. <https://doi.org/10.1155/2020/8361602>

6. Dezutter J. Cognitive and affective aspects of religion in relation to mental health: An exploration in community and clinical samples [PhD thesis]. KU Leuven; 2010. https://kuleuven.limo.libis.be/discovery/fulldisplay?docid=lirias1788445&context=SearchWebhook&vid=32KUL_KU L:Lirias&lang=en&search_scope=lirias_profile&adaptor=SearchWebhook&tab=LIRIAS&query=any,contains,LIRIAS1788445&offset=0

7. Sherman AC, Simonton S. Effects of personal meaning among patients in primary and specialized care: associations with psychosocial and physical outcomes. *Psychol Health.* 2012;27(4):475-490. <https://doi.org/10.1080/08870446.2011.592983>

8. Pinquart M, Silbereisen RK, Fröhlich C. Life goals and purpose in life in cancer patients. *Support Care Cancer.* 2009;17:253-259. <https://doi.org/10.1007/s00520-008-0450-0>

9. Hsieh CC, Ho CC, Lin ZZ, et al. Trait mindfulness and depressive symptoms in non-small cell lung cancer patients: the mediating roles of quality of life and meaning in life. *Psychol Health.* 2021;36(9):1102-1114. <https://doi.org/10.1080/08870446.2020.1825713>

10. Zimmermann FF, Jordan J, Burrell B. Coping with cancer mindfully: a feasibility study of a mindfulness intervention focused on acceptance and meaning in life for adults with advanced cancer. *Eur J Oncol Nurs.* 2020;44:101715. <https://doi.org/10.1016/j.ejon.2019.101715>

11. Zhang JM, Zhang MR, Yang CH, Li Y. The meaning of life according to patients with advanced lung cancer: a qualitative study. *Int J Qual Stud Health Well-being.* 2022;17(1):2028348. <https://doi.org/10.1080/17482631.2022.2028348>

12. Erci B. Meaning in life of patients with cancer. *Palliat Support Care.* Feb 2015;13(1):3-10. <https://doi.org/10.1017/S1478951513000254>

13. Vos J. Cardiovascular disease and meaning in life: a systematic literature review and conceptual model. *Palliat Support Care.* 2021;19(3):367-376. <https://doi.org/10.1017/S1478951520001261>

14. Liu MH, Wang CH, Tung TH, Lee CM, Chiou AF. Care needs, social support and meaning in life in patients after acute heart failure hospitalisation: a longitudinal study. *Eur J Cardiovasc Nurs.* 2021;20(2):106-114. <https://doi.org/10.1177/1474515120945478>

15. Liu MH, Chiou AF, Wang CH, Yu WP, Lin MH. Relationship of symptom stress, care needs, social support, and meaning in life to quality of life in patients with heart failure from the acute to chronic stages: a longitudinal study. *Health Qual Life Outcomes.* 2021;19:252. <https://doi.org/10.1186/s12955-021-01885-8>

16. Barsaei M, Feizollahzadeh H, Jabbarzadeh F, Hasankhani H, Shabanloei R. The meaning of life related to heart disease and its relationship to quality of life after repeat coronary artery angioplasty. *Creat Nurs.* 2020;26(1):48-55. <https://doi.org/10.1891/1078-4535.26.1.48>

17. deRoos-Cassini TA, de St Aubin E, Valvano A, Hastings J, Horn P. Psychological well-being after spinal cord injury: perception of loss and meaning making. *Rehabil Psychol.* 2009;54(3):306-314. <https://doi.org/10.1037/a0016545>

18. Sherman AC, Simonton S, Latif U, Bracy L. Effects of global meaning and illness-specific meaning on health outcomes among breast cancer patients. *J Behav Med.* 2010;33:364-377. <https://doi.org/10.1007/s10865-010-9267-7>

19. Park CL, Edmondson D, Fenster JR, Blank TO. Meaning making and psychological adjustment following cancer: the mediating roles of growth, life meaning, and restored just-world beliefs. *J Consult Clin Psychol.* 2008;76(5):863-875. <https://doi.org/10.1037/a0013348>

20. Yanez B, Edmondson D, Stanton AL, et al. Facets of spirituality as predictors of adjustment to cancer: relative contributions of having faith and finding meaning. *J Consult Clin Psychol.* 2009;77(4):730-741. <https://doi.org/10.1037/a0015820>

21. Vos T, Barber RM, Bell B, et al. Global, regional, and national incidence, prevalence, and years lived with disability for 301 acute and chronic diseases and injuries in 188 countries, 1990-2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet.* 2015;386(9995):743-800. [https://doi.org/10.1016/S0140-6736\(15\)60692-4](https://doi.org/10.1016/S0140-6736(15)60692-4)

22. Drop B, Janiszewska M, Barańska A, Kanecki K, Nitsch-Osuch A, Bogdan M. Satisfaction with life and adaptive reactions in people treated for chronic obstructive pulmonary disease. In: Pokorski, M. (eds) *Clinical Pulmonary Research.* Springer, Cham; 2018: 41-47. https://doi.org/10.1007/5584_2018_242

23. Szymanska-Chabowska A, Juzwiszyn J, Tański W, Świątkowski F, Kobecki J, Chabowski M. The fatigue and quality of life in patients with chronic pulmonary diseases. *Sci Prog.* 2021;104(3). <https://doi.org/10.1177/00368504211044034>
24. Miravittles M, Soriano JB, García-Río F, et al. Prevalence of COPD in Spain: impact of undiagnosed COPD on quality of life and daily life activities. *Thorax.* 2009;64(10):863-868. <https://doi.org/10.1136/thx.2009.115725>
25. Seemungal TAR, Donaldson GC, Paul EA, Bestall JC, Jeffries DJ, Wedzicha JA. Effect of exacerbation on quality of life in patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med.* 1998;157(5):1418-1422. <https://doi.org/10.1164/ajrccm.157.5.9709032>
26. Marx G, Nasse M, Stanze H, Boakye SO, Nauck F, Schneider N. Meaning of living with severe chronic obstructive lung disease: a qualitative study. *BMJ Open.* 2016;6(12):e011555. <https://doi.org/10.1136/bmjopen-2016-011555>
27. Global Initiative for Chronic Obstructive Lung Disease (GOLD). Global strategy for the diagnosis, management, and prevention of COPD, 2023 report. GOLD website. Published 2023. Accessed November 2023. <https://www.goldcopd.org>.
28. Cauty A, Ripley B. *boot: Bootstrap R (S-Plus)* [computer program]. R package version 1.3-28.1. Vienna, Austria: R Foundation for Statistical Computing; 2024. <https://www.r-project.org/foundation/>
29. Davison AC, Hinkley DV. *Bootstrap Methods and Their Applications*. Cambridge University Press; 1997. <https://doi.org/10.1017/CBO9780511802843>
30. R Core Team. *R: A language and environment for statistical computing* [computer program]. Vienna, Austria: R Foundation for Statistical Computing; 2021. <https://www.r-project.org/foundation/>
31. Karasu F, Aylaz R. Evaluation of meaning of life and self-care agency in nursing care given to chronic obstructive pulmonary patients according to health promotion model. *Appl Nurs Res.* 2020;51:151208. <https://doi.org/10.1016/j.apnr.2019.151208>
32. Cramm JM, Nieboer AP. The relationship between self-management abilities, quality of chronic care delivery, and wellbeing among patients with chronic obstructive pulmonary disease in The Netherlands. *Int J Chron Obstruct Pulmon Dis.* 2013;8:209-214. <https://doi.org/10.2147/COPD.S42667>
33. Benzo R, Hoult J, McEvoy C, et al. Promoting chronic obstructive pulmonary disease wellness through remote monitoring and health coaching: a clinical trial. *Ann Am Thorac Soc.* 2022;19(11):1808-1817. <https://doi.org/10.1513/AnnalsATS.202203-214OC>
34. Guerrero-Torrelles M, Monforte-Royo C, Rodríguez-Prat A, Porta-Sales J, Balaguer A. Understanding meaning in life interventions in patients with advanced disease: a systematic review and realist synthesis. *Palliat Med.* 2017;31(9):798-813. <https://doi.org/10.1177/0269216316685235>
35. Harrop E, Noble S, Edwards M, Sivell S, Moore B, Nelson A. Managing, making sense of and finding meaning in advanced illness: a qualitative exploration of the coping and wellbeing experiences of patients with lung cancer. *Sociol Health Illn.* 2017;39(8):1448-1464. <https://doi.org/10.1111/1467-9566.12601>
36. Aktürk Ü, Erci B. The effect of Watson's human caring model on meaning of life and symptom management in cancer patients undergoing chemotherapy. *Res Theory Nurs Pract.* 2018;32(3):255-275. <https://doi.org/10.1891/1541-6577.32.3.255>
37. Winger JG, Ramos K, Steinhauser KE, et al. Enhancing meaning in the face of advanced cancer and pain: Qualitative evaluation of a meaning-centered psychosocial pain management intervention. *Palliat Support Care.* 2020;18(3):263-270. <https://doi.org/10.1017/S1478951520000115>
38. Manco N, Hamby S. A meta-analytic review of interventions that promote meaning in life. *Am J Health Promot.* 2021;35(6):866-873. <https://doi.org/10.1177/0890117121995736>