

Online Supplement

Original Research

Effect of Common Medications on Longitudinal Pectoralis Muscle Area in Smokers

Toru Shirahata, MD^{1,2} Nicholas A. Enzer, BS^{1,3} Victor Castro, BS⁴ Joe Chiles, MD⁵ Merry-Lynn McDonald, PhD^{5,6,7} Bina Choi, MD^{1,3} Alejandro A. Diaz, MD, MPH^{1,3} George R. Washko, MD, MSc^{1,3} Raúl San José Estépar, PhD^{1,8} Samuel Y. Ash, MD, MPH^{9,10} Farbod N. Rahaghi, MD, PhD^{1,3}

¹Applied Chest Imaging Laboratory, Brigham and Women's Hospital, Harvard Medical School, Boston, Massachusetts, United States

²Department of Respiratory Medicine, Saitama Medical University, Saitama, Japan

³Division of Pulmonary and Critical Care Medicine, Department of Medicine, Brigham and Women's Hospital, Harvard Medical School, Boston, Massachusetts, United States

⁴Boston University School of Medicine, Boston, Massachusetts, United States

⁵Division of Pulmonary, Allergy and Critical Care Medicine, Department of Medicine, University of Alabama at Birmingham School of Medicine, Birmingham, Alabama, United States

⁶Department of Genetics, University of Alabama at Birmingham School of Medicine, Birmingham, Alabama, United States

⁷Department of Epidemiology, University of Alabama at Birmingham School of Public Health, Birmingham, Alabama, United States

⁸Department of Radiology, Brigham and Women's Hospital, Harvard Medical School, Boston, Massachusetts, United States

⁹Department of Critical Care Medicine, South Shore Hospital, South Weymouth, Massachusetts, United States

¹⁰Tufts University School of Medicine, Boston, Massachusetts, United States

e-Figure 1. The relationship between longitudinal skeletal muscle density and medication use of propensity matched subjects. (A) delta-PMD with or without statin use, (B) delta-PMD with or without aspirin use. Statin user inhibited PMD decrease (*p=0.041) but no significant difference was found with or without aspirin use(p=0.566). The boxplot shows median delta-PMD and IQR.

