

## Online Supplement

### Brief Report

#### Symptomatic Changes Associated With a Pharmacist-Led COPD Program

Edward C. Portillo, PharmD<sup>1,2</sup> Steven Do, PharmD<sup>1,2,3</sup> Scott Hetzel, MS<sup>4</sup> Jennifer Nguyen<sup>1</sup> Katelynn Cleveland, MS<sup>1</sup> Cara Ray, PhD<sup>5</sup> Jenna Vande Hey<sup>1</sup> Pramit Maskey, MS<sup>6</sup> Rena Steiger-Chadwick, MPH<sup>5</sup> Lucas M. Donovan, MD, MS<sup>7</sup> Dylan Erdelt<sup>1</sup> Sarah Will, PharmD, BCPS<sup>8</sup> Michael Shawn McFarland, PharmD, FCCP, BCACP<sup>8</sup> Heather Ourth, PharmD, BCPS, BCGP, FASHP<sup>8</sup> Michelle Chui, PharmD, PhD<sup>9</sup>

<sup>1</sup>School of Pharmacy, University of Wisconsin-Madison, Madison, Wisconsin, United States

<sup>2</sup>Pharmacy Department, William S. Middleton Veterans Affairs Hospital, Madison, Wisconsin, United States

<sup>3</sup>University of Pittsburgh, Pittsburgh, Pennsylvania, United States

<sup>4</sup>Department of Biostatistics and Medical Informatics, University of Wisconsin-Madison, Madison, Wisconsin, United States

<sup>5</sup>Center for Innovation for Complex Chronic Healthcare, Edward Hines, Jr. Veterans Administration Hospital, Hines, Illinois, United States

<sup>6</sup>Colorado State University, Fort Collins, Colorado, United States

<sup>7</sup>Seattle VA Medical Center, Seattle, Washington, United States

<sup>8</sup>Department of Veterans Affairs Clinical Pharmacy Practice Office, Washington D.C., United States

<sup>9</sup>School of Nursing, University of Wisconsin-Madison, Madison, Wisconsin, United States

**Table 1. Associations between pharmacist interventions and CAT Score Improvement  $\geq 2$**

<b>Pharmacist Intervention, n (%)</b>	<b>CAT Score Improvement <math>\geq 2</math></b>		<b>P-value</b>
	<b>No (N=133)</b>	<b>Yes (N=193)</b>	
<b>Medication Changed - added, removed, or changed a dose of the medication.</b>	70 (53.8%)	138 (71.9%)	0.001
<b>Incorrect technique identified - found the patient to be using the inhaler incorrectly.</b>	31 (23.3%)	31 (16.1%)	0.135
<b>Incorrect adherence identified - found the patient to be nonadherent to a COPD Medication.</b>	46 (34.6%)	71 (36.8%)	0.772
<b>Service referral – referred patients to at least one service (such as telehealth, pulmonary clinic, COPD Class, etc.) to support COPD Management.</b>	59 (45.4%)	92 (47.9%)	0.739
<b>COPD Action Plan - completed a physical action plan indicating to the veteran what to do in the event of worsening symptoms.</b>	89 (68.5%)	147 (76.6%)	0.138

Reported p-values are based on chi-square tests of independence assessing whether the proportion of those receiving each intervention significantly differed between those who experienced clinically meaningful improvement and those who did not,  $\alpha = 0.05$ .

Note: The COPD CARE note template captured medication changes of addition, removal, or dose change and incorrect adherence identification without specifying which medications were involved.

COPD=Chronic Obstructive Pulmonary Disease; CAT=COPD Assessment Test