## **Online Supplement**

## Characterizing COPD Symptom Variability in the Stable State Utilizing the Evaluating Respiratory Symptoms in COPD Questionnaire

Jamuna K. Krishnan, MD<sup>1\*</sup> Kayley M. Ancy<sup>2\*</sup> Clara Oromendia<sup>3</sup>, Katherine L. Hoffman, PhD<sup>3</sup> Imaani Easthausen<sup>3</sup> Nancy K. Leidy, PhD<sup>4</sup> MeiLan K. Han, MD, MS<sup>5</sup> Russell P. Bowler, MD, PhD<sup>6</sup> Stephanie A. Christenson, MD<sup>7</sup> David J. Couper, PhD<sup>8</sup> Gerard J. Criner, MD<sup>9</sup> Jeffrey L. Curtis, MD<sup>5,10</sup> Mark T. Dransfield, MD<sup>11</sup> Nadia N. Hansel, MD, MPH<sup>12</sup> Anand S. Iyer<sup>11</sup>, Robert Paine III, MD<sup>13</sup> Stephen P. Peters, MD, PhD<sup>14</sup> Jadwiga A. Wedzicha, MD<sup>15</sup> Prescott G. Woodruff, MD, MPH<sup>7</sup> Karla V. Ballman<sup>3</sup> Fernando J. Martinez, MD, MS<sup>1</sup> for the SPIROMICS Investigators

<sup>\*</sup>Co-first authors, both authors contributed equally to the work.

<sup>&</sup>lt;sup>1</sup> Pulmonary and Critical Care Medicine, Weill Cornell Medicine, New York, New York, United States

<sup>&</sup>lt;sup>2</sup> Department of Medicine, Weill Cornell Medicine, New York, New York, United States

<sup>&</sup>lt;sup>3</sup> Division of Biostatistics, Department of Population Health Sciences, Weill Cornell Medical College, New York, New York, United States

<sup>&</sup>lt;sup>4</sup> Evidera, Bethesda, Maryland, United States

<sup>&</sup>lt;sup>5</sup> Internal Medicine, University of Michigan Health System, Ann Arbor, Michigan, United States

<sup>&</sup>lt;sup>6</sup> Department of Medicine, National Jewish Medical and Research Center, Denver, Colorado, United States

<sup>&</sup>lt;sup>7</sup> Pulmonary and Critical Care, University of California San Francisco, San Francisco, California, United States

<sup>&</sup>lt;sup>8</sup> University of North Carolina at Chapel Hill, Chapel Hill, North Carolina, United States

<sup>&</sup>lt;sup>9</sup> Pulmonary and Critical Care Medicine, Temple University Hospital, Philadelphia, Pennsylvania, United States

<sup>&</sup>lt;sup>10</sup> Medical Service, VA Ann Arbor Healthcare System, Ann Arbor, Michigan, United States

<sup>&</sup>lt;sup>11</sup> Pulmonary, Allergy and Critical Care, University of Alabama at Birmingham, Birmingham, Alabama, United States

<sup>&</sup>lt;sup>12</sup> Division of Pulmonary and Critical Care Medicine, Johns Hopkins University, Baltimore Maryland, United States

<sup>&</sup>lt;sup>13</sup> Division of Pulmonary and Critical Care Medicine, Department of Veterans Affairs Medical Center, University of Utah, Salt Lake City, Utah, United States

<sup>&</sup>lt;sup>14</sup> Section on Pulmonary, Critical Care, Allergy and Immunological Diseases, Wake Forest School of Medicine Medical Center, Winston-Salem, North Carolina, United States

<sup>&</sup>lt;sup>15</sup> National Heart and Lung Institute, Imperial College London, United Kingdom

## **Supplementary Tables**

Table E1. Comparison of included and excluded subjects in the symptom variability analysis

Characteristic	Included	Excluded	Overall
N	140 [68.3%]	65 [31.7%]	205
Age (years)	63.3 ±8.4	$63.3 \pm 7.6$	63.3 ±8.1
Female sex, n [%]	74 [52.9%]	29 [44.6%]	103 [50.2%]
BMI* (kg/m²)	27.5 ±5.5	27.2 ±5.5	27.4 ±5.5
Six-minute walk distance (10m)	36.2 ±12.3	38.7 ±12.5	37.0 ±12.4
FEV <sub>1</sub> <sup>†</sup> % predicted	61.5 ±21.9	$65.0 \pm 21.9$	62.6 ±21.8
BODE <sup>‡</sup> Index	2.3 ±2.2	$1.8 \pm 2.1$	2.1 ±2.2
MMRC§ dyspnea scale	$1.4 \pm 1.1$	$1.0\pm1.0$	1.3 ±1.0
PSQ <sup>  </sup> total score	$6.9 \pm 3.8$	$7.2 \pm 4.6$	$7.0 \pm 3.9$
SGRQ** total score	$42.9 \pm 19.9$	$36.5 \pm 19.9$	41.1 ±21.0
HADS <sup>††</sup> anxiety score	6.2 ±4.0	$5.6 \pm 4.0$	6.0 ±4.0
HADS <sup>††</sup> depression score	$4.9 \pm 3.7$	$4.5 \pm 3.3$	$4.8 \pm 3.5$
Current smoker	47 [33.8%]	27 [41.5%]	74 [36.3%]
GOLD <sup>‡‡</sup> stage, n [%]			
Stage I	11 [7.9%]	4 [6.3%]	15 [7.4%]
Stage II	31 [22.3%]	14 [21.5%]	45 [22.1%]
Stage III	67 [48.2%]	29 [44.6%]	96 [47.1%]
Stage IV	30 [21.6%]	18 [23.8%]	48 [23.5%]
Unavailable	1 [0.7%]	0 [3.1%]	1 [0.5%]
EXACT§§ mean	35.2 ±11.0	33.4 ±13.1	34.7 ±11.5
E-RS <sup>∆</sup> mean	11.4 ±6.6	9.9 ±7.5	11.0 ±6.9
E-RS <sup>∆</sup> subscales			
E-RS breathlessness	$5.6 \pm 3.8$	$4.9 \pm 4.2$	5.4 ±4.0
E-RS cough/sputum	3.2 ±1.9	2.9 ±1.9	3.1 ±1.9
E-RS chest	2.6 ±2.0	2.2 ±2.1	2.4 ±2.0
WS-SD <sup>∆∆</sup> at baseline	$3.04 \pm 1.52$	$2.55 \pm 1.74$	$2.89 \pm 1.60$
HCRU <sup>^</sup> Exacerbations	18 [12.9%]	3[4.6%]	21[10.2%]
EXACT <sup>§§</sup> event within 4 weeks of follow-up (1 or more event), n [%]	45 [32.1%]	5 [7.7%]	50 [24.4%]

\*BMI = Body mass index; †FEV1 = Forced expiratory volume in one second; †BODE = Body mass index, airflow obstruction, dyspnea and exercise capacity; MMRC = Modified Medical Research Council; PSQ = Pittsburgh Sleep Quality Index; \*\*SGRQ = St. George Respiratory Questionnaire; †HADS = hospital anxiety and depression scale; †GOLD = Global Initiative For Chronic Obstructive Lung Disease; <math>EXACT = EXACT = Exacerbations of Chronic Pulmonary Disease Tool; EXSCT = EXACT = Evaluating Respiratory Symptoms in COPD Tool; EXSCT = EXACT = Within-subject standard deviation; EXSCT = EXACT = Evaluation exacerbations in the 52 weeks following Exacerbation sub study enrollment.

Table E2. Pair wise differences in weekly WS-SD\* among 130 subjects with sufficient data in 12 weeks

Start Week	End Week											
	1	2	3	4	5	6	7	8	9	10	11	12
0	0.07	-0.05	-0.34	-0.45	-0.4	-0.6	-0.78	-0.57	-0.67	-0.48	-0.62	-0.64
	p <sup>†</sup> =0.71	p=0.72	p=0.025	p=0.006	p=0.007	p<0.001	p<0.001	p<0.001	p<0.001	p=0.007	p<0.001	p<0.001
1		-0.12	-0.4	-0.52	-0.46	-0.66	-0.85	-0.64	-0.74	-0.55	-0.69	-0.7
		p=0.53	p=0.035	p=0.007	p=0.008	p<0.001	p<0.001	p=0.001	p<0.001	p=0.006	p<0.001	p<0.001
2			-0.28	-0.39	-0.34	-0.54	-0.73	-0.52	-0.62	-0.42	-0.57	-0.58
			p=0.039	p=0.02	p=0.014	p<0.001	p<0.001	p<0.001	p<0.001	p=0.014	p<0.001	p<0.001
3				-0.11	-0.06	-0.26	-0.45	-0.23	-0.33	-0.14	-0.28	-0.3
				p=0.48	p=0.66	p=0.075	p=0.003	p=0.095	p=0.032	p=0.41	p=0.087	p=0.056
4					0.05	-0.15	-0.34	-0.12	-0.22	-0.03	-0.17	-0.19
					p=0.69	p=0.30	p=0.022	p=0.38	p=0.11	p=0.86	p=0.28	p=0.24
5						-0.2	-0.39	-0.17	-0.27	-0.08	-0.22	-0.24
						p=0.078	p=0.03	p=0.17	p=0.03	p=0.60	p=0.11	p=0.095
6							-0.19	0.02	-0.07	0.12	-0.03	-0.04
							p=0.14	p=0.85	p=0.57	p=0.44	p=0.85	p=0.77
7								0.21	0.11	0.31	0.16	0.15
								p=0.11	p=0.42 -0.1	p=0.068 0.09	p=0.26 -0.05	p=0.28 -0.07
8									p=0.42	p=0.55	p=0.03	p=0.66
									p=0.42	0.19	0.05	0.03
9										p=0.20	p=0.73	p=0.82
10										P 0.20	-0.14	-0.16
											p=0.37	p=0.31
											P 0.57	-0.02
11												p=0.91

<sup>\*</sup>WS-SD = within subject standard deviation. †all p-values indicate paired t-tests, bold indicates statistical significance.

## SUPPLEMENTARY FIGURES

**Figure E1.** Example of one subject's E-RS scores throughout the study with standard deviation (blue dashed line) decreasing throughout the weeks. Day -7 to day 0 represents the baseline week. Mean E-RS score (red line) demonstrates more stability during the same time period. E-RS = Evaluating Respiratory Symptoms in COPD Tool. WS-SD = within subject standard deviation

**Figure E2.** Mean WS-SD of subjects over twelve weeks of follow-up after baseline week. Grey lines indicate each individual subject's WS-SD trajectory over the 13 weeks. WS-SD = within subject standard deviation

**Figure E3.** Example of two subject's daily EXACT scores leading to the onset and recovery of an EXACT event.

Figure E1




Figure E2




Figure E3

