

Original Research**Fall Risk and Medication Use Near End of Life Among Adults With Chronic Obstructive Pulmonary Disease**

Cara L. McDermott, PharmD, PhD^{1,2} Laura C. Feemster, MD, MS^{3,4} Ruth A. Engelberg, PhD³
Laura J. Spece, MD, MS^{3,4} Lucas M. Donovan, MD, MS^{3,4} J. Randall Curtis, MD, MPH^{3†}

¹Division of Geriatrics, Department of Medicine, Duke University School of Medicine, Durham, North Carolina, United States

²Department of Population Health Sciences, Duke University School of Medicine, Durham, North Carolina, United States

³Division of Pulmonary, Critical Care and Sleep Medicine, University of Washington, Seattle, Washington, United States

⁴Health Systems Research, Department of Veterans Affairs Puget Sound Health Care System, Seattle, Washington, United States

†deceased

Address correspondence to:

Cara L. McDermott, PharmD, PhD
Box 3003, DUMC
Division of Geriatrics
Durham, NC 27710
Phone: (919) 660-8386
Email: cara.mcdermott@duke.edu

Running Head: Falls and Medications Among Patients With COPD

Keywords: COPD; falls; medication; end of life; care delivery

Abbreviations: COPD=chronic obstructive pulmonary disease, FRID=fall risk increasing drug, OR=odds ratio, CI=confidence interval, EHR=electronic health record, ICD=International Classification of Disease, SSRI=selective serotonin reuptake inhibitor, SNRI=serotonin norepinephrine reuptake inhibitor, SD=standard deviation

Funding Support: Grant support provided by NIH/NHLBI K23 HL159239 (CLM), Cambia Foundation (RAE, JRC), American Lung Association (LCF), PCORI (LCF, LD), and the Department of Veterans Affairs (LCF, LJS, LD).

Date of Acceptance: October 21, 2024 | **Publication Online Date:** October 25, 2024

Citation: McDermott CL, Feemster LC, Engelberg RA, Spece LJ, Donovan LM, Curtis JR. Fall risk and medication use near end of life among adults with chronic obstructive pulmonary disease. *Chronic Obstr Pulm Dis.* 2024; Published online October 25, 2024.

This article has an [online supplement](#).

Abstract**Background:**

Falls are frequent among people with chronic obstructive pulmonary disease (COPD) and associated with increased morbidity, mortality, and healthcare costs. Understanding modifiable medication factors that contribute to fall risk is an important step to developing fall prevention strategies for this highly susceptible group.

Methods:

This is a retrospective cohort study using electronic health record data from a single health system linked to Washington State death certificates of adults ages 40 or older who died between 2014-2018 with COPD. We identified demographics, comorbidities, fall-risk increasing drug (FRID) burden, and the occurrence of injurious falls within the 2 years prior to the date of death. We defined injurious falls using published algorithms of International Classification of Disease codes.

Results:

Of 8204 decedents with COPD, 2454 (30%) had an injurious fall in the two years before death, and FRID use was common among 65%. A higher percentage of patients with falls received prescriptions for anticonvulsants (35% vs 26%), antipsychotics (24% vs 13%), atypical antidepressants (28% vs 19%), tricyclic antidepressants (10% vs 5%) versus those without a fall. In multivariable logistic regression, after adjusting for confounders, FRID burden was associated with greater odds of injurious fall (odds ratio (OR) 1.07 (95% confidence interval (CI) 1.04-1.09).

Conclusion:

Our findings highlight an opportunity for collaboration between pharmacists, pulmonologists, and patients to develop new processes to potentially deprescribe and optimize the use of FRIDs among patients with COPD to increase safety.

Introduction

Patients with chronic obstructive pulmonary disease (COPD) are significantly more likely to experience a fall compared to patients without COPD,(1-4) with increased fall risk due to increased frailty,(5) impaired gait,(2) and tripping hazards from oxygen tubing.(6) Falls and resulting injuries such as fractures can result in emergency department visits or hospitalizations, which lower patient quality of life and increase healthcare costs.(7-10) Despite COPD being the sixth most common cause of death in the United States, little is known about falls among these patients near the end of life.(11)

One modifiable risk factor for falls is medication use, including fall-risk increasing drugs (FRIDs) such as benzodiazepines, sedative-hypnotics, antihypertensives, and other psychoactive agents.(12) Patients with COPD have a high prevalence of comorbidities for which FRIDs are often used as treatment, including anxiety, depression, diabetes, and hypertension. The burden of polypharmacy also increases as patients' function declines, and they receive additional medications for insomnia and pain.(13-15)

While these medications are intended to alleviate patient symptoms, they also increase the risk of injurious falls.(16-20) Reducing FRID burden represents an opportunity to improve patient outcomes. While previous work has found an association between polypharmacy and increased fall risk, (21) FRID burden among patients with COPD, and the relationship between FRID burden and fall risk has not been previously described. FRID use among patients with COPD is of particular concern given the large number of FRID indications (e.g. anxiety, insomnia, dyspnea) among patients with COPD.

We conducted a retrospective cohort study in the University of Washington (UW) Medicine system, which is the largest public health system in the Puget Sound area. It includes an academic medical center, a county safety-net hospital, and two community hospitals, plus outpatient clinics. With these data, we evaluated the association between FRID use and injurious falls in the last two years of life among decedents with COPD. As FRID exposure by class is not well described among patients with COPD, we describe the frequency of FRID prescriptions for different classes of FRIDs in this cohort (e.g. benzodiazepines, sedative hypnotics, tricyclic antidepressants), comparing the frequency of FRID prescriptions among patients with COPD who experienced an injurious fall to patients with COPD without an injurious fall. We included

anti-convulsants for patients without comorbid seizure disorder. Our hypothesis is that among patients with COPD, injurious falls are associated with FRID burden.

Methods

Study Design and Population

For this study we used a linked database between Washington State death certificates to the electronic health record (EHR) of the UW Medicine system to evaluate patterns of care for adults with serious illness. The goal of study was to identify factors associated with an increased risk of injurious falls for adults with COPD who died from any cause between 2014-2018. COPD was defined using International Classification of Disease (ICD) codes, specifically ICD-9-CM 491.xx, 492.xx, or 496, and ICD-10-CM J41, J42, J43, J44. Patients had to have at least two codes assigned to be classified as having COPD.

Exposure and Outcome

Primary exposure of interest: We identified the presence or absence of prescriptions for FRIDs in the EHR and grouped medications by class. (18, 22, 23) These medication classes included anticonvulsants, antihypertensives, antipsychotics, antispasmodics, atypical antidepressants (e.g. bupropion, mirtazapine), benzodiazepines, opioids, sedative-hypnotics, selective serotonin reuptake inhibitors (SSRIs), serotonin norepinephrine reuptake inhibitors (SNRIs), plus tricyclic and tetracyclic antidepressants. Medications were identified using brand or generic name in the EHR (full list available in Supplemental Table 1). Patients with a prescription for a FRID of any duration were categorized as exposed to the medication, and we reported the total number of FRID prescriptions per patient by class (i.e. “FRID burden”).

Primary Outcome: We identified an injurious fall using ICD-9 code V15.88 and ICD-10 code Z91.81.(24) Our outcome is documentation of an injurious fall requiring medical attention, drawn from previously published algorithms and defined as emergency department visits, hospitalizations, and outpatient visits with a fall-related diagnosis and with a code for: brain injury; contusion; dislocation of the hip, jaw, knee, shoulder, wrist; non-pathological fracture of the skull, face, spine, clavicle, ribs, humerus, wrist, pelvis, hip, fibula, tibia, foot or ankle.(24, 25) We excluded visits for a motor vehicle collision.(26) The full list of ICD codes is available in Supplemental Table 2.

Adjustment confounders

The following confounders were selected based on prior data suggesting an association with risks of fall: (27-30) comorbidities, race (previously self-reported), level of education (completed some college or higher), and marital status (married or not married at the time of death) from the Washington State death certificates; age, sex, and insurance status (private, Medicare, Medicaid, other) were identified from the EHR.

Using an adaptation of the Dartmouth Atlas criteria for serious illness, we identified comorbidities and fall history from the EHR using ICD codes versions 9 or 10, using ICD-10 codes for diagnoses after 2015.(31) In addition, all patients could have an additional comorbidity as defined by the Dartmouth Atlas (31): advanced renal disease, coronary artery disease, dementia, diabetes with end organ damage, end-stage liver disease, heart failure, pulmonary disease other than COPD (e.g. idiopathic pulmonary fibrosis), peripheral vascular disease, poor prognosis cancer (e.g. metastatic solid tumor or aggressive hematologic malignancy such as acute myeloid leukemia).(31) Together, these conditions comprise the most common causes of death in America, excluding suicide, accidents and infections.(32)

We refined and updated these codes through continuous quality improvement, using the general equivalency maps provided by the Centers for Medicare and Medicaid Services(33) in addition to manual review by two clinicians to resolve any potential discrepancies in mapping codes to conditions. Each comorbidity was assessed separately rather than using a summed comorbidity score. Since comorbidity prevalence varies among patients with COPD, this afforded us the opportunity to illustrate the comorbidity patterns in this population. We used ICD codes to identify the presence of comorbidities. Unfortunately, ICD codes do not indicate condition severity, thus we assigned all comorbidities equal weight.

Statistical analysis

We describe the study population by comorbidities including fall history, demographics, and FRID prescriptions for those with and without an injurious fall, using a t-test to examine differences in FRID prescription type between groups at 95% significance. Using multivariable logistic regression in a single model, we examined the association of FRID burden with the outcome of injurious fall, dichotomized as a binary variable of 0 (none) or 1 (1 fall or more),

while adjusting for demographic characteristics, types of comorbidities, fall history, and FRID burden, defined as number of FRID prescriptions for different FRID classes. The University of Washington Institutional Review Board approved this study and issued a waiver of consent to access decedent records in accordance with Washington State law. We performed all analyses using Stata version 16.0 (College Station, TX).

Results

We identified 8,204 decedents with COPD, and of these patients, 2,454 (30%) had an injurious fall. Patients with COPD and an injurious fall were more likely to be Black, unmarried, and enrolled in Medicaid, while those without an injurious fall more frequently had private insurance and at least some college-level education. (Table 1) A history of falling was markedly higher among patients experiencing an injurious fall compared to those without a fall (23% vs 4%). Patients with COPD who had an injurious fall also had more comorbidities compared to those with COPD who did not fall (Table 2). The average number of comorbidities was 2.81 (standard deviation (SD) 1.80) among those with an injurious fall versus 2.46 (SD 1.56) for those without a fall. The only comorbidity that occurred more frequently among those without an injurious fall was cancer (26% among patients who fell vs 35% with no fall).

In multivariate logistic regression, FRID prescription was associated with a higher risk of injurious falls (odds ratio (OR) 1.07 (95% confidence interval (CI) 1.04-1.09). FRID burden was higher among patients with an injurious fall, with a mean of 3.47 prescriptions across FRID classes (SD 2.66) whereas those without an injurious fall had an average FRID burden of 2.85 FRIDs (SD 2.24) (Table 3). Across both groups, the most prescribed FRID classes were antihypertensives, opioids and benzodiazepines. Comparing patients with an injurious fall to those without, the largest differences in prescriptions were noted for anticonvulsants (35% vs 26%), antipsychotics (24% vs 13%), atypical antidepressants (28% vs 19%), tricyclic antidepressants (10% vs 5%) and SSRIs (27% vs 20%). Across all medication classes, FRID use was statistically higher ($p < 0.05$) among those with an injurious fall compared to those without a fall, except for antihypertensives ($p = 0.52$).

Discussion

We found that FRID burden was associated with a higher risk of falls, after adjusting for demographic and clinical conditions. With this analysis we provide additional information on which medication classes could be targeted by interventions to mitigate fall risk and improve patient outcomes.

In situations of polypharmacy, it can be difficult to decide which medications should be prioritized for dose reduction or discontinuation. We note that the most prescribed FRID class among all patients with COPD was antihypertensives. While the difference between the two groups with falls was not statistically significant, this still may be an area to consider deprescribing and/or dose reductions for patients with COPD, especially for those with episodes of orthostatic hypotension, well-controlled blood pressure, or limited life expectancy. In a previous study of older adults that explored ways to address medication-related symptoms, patients with COPD were more likely to prioritize reduction of fall injuries compared to reduction of cardiovascular events.(34)

The second and third most common FRID classes in our analysis were opioids and benzodiazepines. These medications can be prescribed as palliation to relieve breathlessness or anxiety among patients with COPD.(35, 36) While such medications are appropriate in this context, additional fall mitigation options should be considered. We noted that patients with falls were more likely to receive anticonvulsants, antipsychotics, atypical antidepressants, tricyclic antidepressants, and SSRIs. These medications are often used for pain, insomnia, and depression; nonpharmacological options could perhaps reduce some medication use and thus lower fall risk.

Finally, dementia was a more common comorbidity among those with a fall compared to those who did not fall. This finding aligns with other studies that have found patients living with dementia are two to three times more likely to fall compared to patients without dementia,(37, 38) since gait control is negatively impacted when cognition is reduced. When possible, non-pharmacological alternatives to FRIDs, or using the lowest FRID dose possible for the shortest duration should be encouraged for patients living with both COPD and dementia.

There are limitations to this study. First, patients may have received care for falls in other healthcare systems, but these data were not available to us as we relied on one EHR, a known limitation of EHR-based studies.(39) We relied on ICD-9 and ICD-10 coding to determine the presence of comorbidities but these codes do not provide information as to condition severity, which may confound fall risk. We had access to medication orders rather than pharmacy claims

that implied a prescription was filled. It is possible that patients did not take the prescribed medications or did not fill the prescriptions; additionally, we have no information on over-the-counter FRIDs such as diphenhydramine that patients may have taken. Finally, we do not have information about COPD severity/exacerbation history or home oxygen use, both of which are associated with falls.

Medication management is an area where pharmacist collaborations with pulmonologists and primary care providers can personalize care to reduce fall risk when engaging in shared decision making and fall risk mitigation. Our study suggests that additional screening regarding medication use and collaboration across disciplines (e.g. pulmonary, primary care, cardiology, palliative care) may help mitigate fall risk while managing symptoms and comorbidities.

Declaration of Interest

- 1) Cara L. McDermott reports grant funding from NIH/NHLBI. She is a consultant for the non-profit Biologics and Biosimilars Collective Intelligence Consortium and received honoraria from the University of Utah in 2023 for speaking at two departmental seminars.
- 2) Laura C. Feemster received grant support from PCORI, the American Lung Association, the Department of Veterans Affairs Health Systems Research, and NIH/NHLBI. She has received travel funding from the American Thoracic Society to attend leadership meetings, and consultant fees from the Society of Hospital Medicine and the National Quality Assurance Committee.
- 3) Ruth A. Engelberg received funding support from the Cambia Health Foundation.
- 4) Laura J. Spece reports grant support from NIH/NHLBI.
- 5) Lucas Donovan reports grant support from the Department of Veterans Affairs Health Systems Research and PCORI, plus an honorarium from the American Academy of Sleep Medicine as a CME speaker for the 2023 conference.
- 6) J. Randall Curtis received funding support from the Cambia Health Foundation.

Acknowledgements

We appreciate the help of James Sibley with the creation of the database used for analysis.

Author participation and contributions:

- 1) Cara L. McDermott conceptualized the project, conducted formal analysis, obtained funding, conducted the investigation, decided upon and executed study methodology, administered the project, wrote the original draft, and reviewed/edited the final manuscript version.
- 2) Laura C. Feemster helped contextualize results of the formal analysis, contributed to study design/ methodology, co-wrote the original draft, and reviewed/edited the final manuscript version.
- 3) Ruth A. Engelberg helped conceptualize the project, curated data, contributed to formal analyses, obtained funding, co-conducted the investigation, decided upon and executed study methodology, co-administered the project, co-wrote the original draft, and reviewed/edited the final manuscript version.
- 4) Laura J. Spece contributed to formal analysis, contributed to study design/ methodology, co-wrote the original draft, and reviewed/edited the final manuscript version.
- 5) Lucas Donovan contributed to study design/ methodology, co-wrote the original draft, and reviewed/edited the final manuscript version.
- 6) J. Randall Curtis helped conceptualize the project, curated data, obtained funding, co-conducted the investigation, contributed to study design, co-wrote the original draft, and reviewed/edited the final manuscript version.

References

1. Gazzotti MR, Roco CM, Pradella CO, Nascimento OA, Porto EF, Adas M, et al. Frequency of Osteoporosis and Vertebral Fractures in Chronic Obstructive Pulmonary Disease (COPD) Patients. *Arch Bronconeumol*. 2019;55(5):252-7. doi: 10.1016/j.arbres.2018.10.010. PubMed PMID: 30541669.
2. Hakamy A, Bolton CE, Gibson JE, McKeever TM. Risk of fall in patients with COPD. *Thorax*. 2018;73(11):1079-80. doi: 10.1136/thoraxjnl-2017-211008. PubMed PMID: 29563161.
3. Pereira A, Xavier RF, Lopes AC, da Silva C, Oliveira CC, Fernandes FLA, et al. The Mini-Balance Evaluation System Test Can Predict Falls in Clinically Stable Outpatients With COPD: A 12-MO PROSPECTIVE COHORT STUDY. *J Cardiopulm Rehabil Prev*. 2019;39(6):391-6. doi: 10.1097/HCR.0000000000000427. PubMed PMID: 31689266.
4. Oliveira CC, Lee AL, McGinley J, Thompson M, Irving LB, Anderson GP, et al. Falls by individuals with chronic obstructive pulmonary disease: a preliminary 12-month prospective cohort study. *Respirology*. 2015;20(7):1096-101. Epub 20150723. doi: 10.1111/resp.12600. PubMed PMID: 26206432.
5. Gephine S, Mucci P, Grosbois JM, Maltais F, Saey D. Physical Frailty in COPD Patients with Chronic Respiratory Failure. *Int J Chron Obstruct Pulmon Dis*. 2021;16:1381-92. Epub 2021/05/29. doi: 10.2147/COPD.S295885. PubMed PMID: 34045852; PubMed Central PMCID: PMC8144849.
6. Wiener RS, Ouellette DR, Diamond E, Fan VS, Maurer JR, Mularski RA, et al. An official American Thoracic Society/American College of Chest Physicians policy statement: the Choosing Wisely top five list in adult pulmonary medicine. *Chest*. 2014;145(6):1383-91. Epub 2014/06/04. doi: 10.1378/chest.14-0670. PubMed PMID: 24889436; PubMed Central PMCID: PMC4694177.
7. Choi NG, Choi BY, DiNitto DM, Marti CN, Kunik ME. Fall-related emergency department visits and hospitalizations among community-dwelling older adults: examination of health problems and injury characteristics. *BMC Geriatr*. 2019;19(1):303. doi: 10.1186/s12877-019-1329-2. PubMed PMID: 31711437; PubMed Central PMCID: PMC6849272.
8. Marchini L, Allareddy V. Epidemiology of facial fractures among older adults: A retrospective analysis of a nationwide emergency department database. *Dent Traumatol*. 2019;35(2):109-14. doi: 10.1111/edt.12459. PubMed PMID: 30548143.
9. Kragh Ekstam A, Elmstahl S. Do fall-risk-increasing drugs have an impact on mortality in older hip fracture patients? A population-based cohort study. *Clin Interv Aging*. 2016;11:489-96. doi: 10.2147/CIA.S101832. PubMed PMID: 27199553; PubMed Central PMCID: PMC4857759.
10. Zia A, Kamaruzzaman SB, Tan MP. The consumption of two or more fall risk-increasing drugs rather than polypharmacy is associated with falls. *Geriatr Gerontol Int*. 2017;17(3):463-70. doi: 10.1111/ggi.12741. PubMed PMID: 26822931.
11. Xu J, Murphy SL, Kochanek KD, Arias E. Mortality in the United States, 2021. *NCHS Data Brief*. 2022(456):1-8. PubMed PMID: 36598387.
12. By the American Geriatrics Society Beers Criteria Update Expert P. American Geriatrics Society 2023 updated AGS Beers Criteria(R) for potentially inappropriate medication use in older adults. *J Am Geriatr Soc*. 2023;71(7):2052-81. Epub 20230504. doi: 10.1111/jgs.18372. PubMed PMID: 37139824.
13. Pumar MI, Gray CR, Walsh JR, Yang IA, Rolls TA, Ward DL. Anxiety and depression-Important psychological comorbidities of COPD. *J Thorac Dis*. 2014;6(11):1615-31. Epub 2014/12/06. doi: 10.3978/j.issn.2072-1439.2014.09.28. PubMed PMID: 25478202; PubMed Central PMCID: PMC4255157.
14. Weiss A, Porter S, Rozenberg D, O'Connor E, Lee T, Balter M, et al. Chronic Obstructive Pulmonary Disease: A palliative medicine review of the disease, its therapies and drug interactions. *J Pain Symptom Manage*. 2020. doi: 10.1016/j.jpainsymman.2020.01.009. PubMed PMID: 32004618.
15. Duncan I, Maxwell TL, Huynh N, Todd M. Polypharmacy, Medication Possession, and Deprescribing of Potentially Non-Beneficial Drugs in Hospice Patients. *Am J Hosp Palliat Care*. 2020;37(12):1076-85. Epub 2020/07/15. doi: 10.1177/1049909120939091. PubMed PMID: 32662276.
16. Baillargeon J, Singh G, Kuo YF, Raji MA, Westra J, Sharma G. Association of Opioid and Benzodiazepine Use with Adverse Respiratory Events in Older Adults with Chronic Obstructive Pulmonary Disease. *Ann Am Thorac Soc*. 2019;16(10):1245-51. doi: 10.1513/AnnalsATS.201901-024OC. PubMed PMID: 31104504; PubMed Central PMCID: PMC6812171.

17. Donovan LM, Malte CA, Spece LJ, Griffith MF, Feemster LC, Zeliadt SB, et al. Center Predictors of Long-Term Benzodiazepine Use in Chronic Obstructive Pulmonary Disease and Post-traumatic Stress Disorder. *Ann Am Thorac Soc*. 2019;16(9):1151-7. doi: 10.1513/AnnalsATS.201901-048OC. PubMed PMID: 31113231; PubMed Central PMCID: PMC6812159.
18. Seppala LJ, Wermelink A, de Vries M, Ploegmakers KJ, van de Glind EMM, Daams JG, et al. Fall-Risk-Increasing Drugs: A Systematic Review and Meta-Analysis: II. Psychotropics. *J Am Med Dir Assoc*. 2018;19(4):371 e11- e17. Epub 2018/02/07. doi: 10.1016/j.jamda.2017.12.098. PubMed PMID: 29402652.
19. Hart LA, Marcum ZA, Gray SL, Walker RL, Crane PK, Larson EB. The Association Between Central Nervous System-Active Medication Use and Fall-Related Injury in Community-Dwelling Older Adults with Dementia. *Pharmacotherapy*. 2019;39(5):530-43. doi: 10.1002/phar.2244. PubMed PMID: 30861179; PubMed Central PMCID: PMC6983303.
20. Aspinall SL, Springer SP, Zhao X, Cunningham FE, Thorpe CT, Semla TP, et al. Central Nervous System Medication Burden and Risk of Recurrent Serious Falls and Hip Fractures in Veterans Affairs Nursing Home Residents. *J Am Geriatr Soc*. 2019;67(1):74-80. doi: 10.1111/jgs.15603. PubMed PMID: 30306541.
21. Roig M, Eng JJ, MacIntyre DL, Road JD, FitzGerald JM, Burns J, et al. Falls in people with chronic obstructive pulmonary disease: an observational cohort study. *Respir Med*. 2011;105(3):461-9. doi: 10.1016/j.rmed.2010.08.015. PubMed PMID: 20869227; PubMed Central PMCID: PMC3350813.
22. de Vries M, Seppala LJ, Daams JG, van de Glind EMM, Masud T, van der Velde N, et al. Fall-Risk-Increasing Drugs: A Systematic Review and Meta-Analysis: I. Cardiovascular Drugs. *J Am Med Dir Assoc*. 2018;19(4):371 e1- e9. Epub 2018/02/06. doi: 10.1016/j.jamda.2017.12.013. PubMed PMID: 29396189.
23. Seppala LJ, van de Glind EMM, Daams JG, Ploegmakers KJ, de Vries M, Wermelink A, et al. Fall-Risk-Increasing Drugs: A Systematic Review and Meta-analysis: III. Others. *J Am Med Dir Assoc*. 2018;19(4):372 e1- e8. Epub 2018/02/07. doi: 10.1016/j.jamda.2017.12.099. PubMed PMID: 29402646.
24. Hoffman GJ, Hays RD, Shapiro MF, Wallace SP, Ettner SL. Claims-based Identification Methods and the Cost of Fall-related Injuries Among US Older Adults. *Med Care*. 2016;54(7):664-71. doi: 10.1097/MLR.0000000000000531. PubMed PMID: 27057747; PubMed Central PMCID: PMC4907826.
25. Min L, Tinetti M, Langa KM, Ha J, Alexander N, Hoffman GJ. Measurement of Fall Injury With Health Care System Data and Assessment of Inclusiveness and Validity of Measurement Models. *JAMA Netw Open*. 2019;2(8):e199679. Epub 20190802. doi: 10.1001/jamanetworkopen.2019.9679. PubMed PMID: 31433480; PubMed Central PMCID: PMC6707014.
26. Green AR, Reifler LM, Bayliss EA, Weffald LA, Boyd CM. Drugs Contributing to Anticholinergic Burden and Risk of Fall or Fall-Related Injury among Older Adults with Mild Cognitive Impairment, Dementia and Multiple Chronic Conditions: A Retrospective Cohort Study. *Drugs Aging*. 2019;36(3):289-97. doi: 10.1007/s40266-018-00630-z. PubMed PMID: 30652263; PubMed Central PMCID: PMC6386184.
27. Oliveira CC, Annoni R, Lee AL, McGinley J, Irving LB, Denehy L. Falls prevalence and risk factors in people with chronic obstructive pulmonary disease: A systematic review. *Respir Med*. 2021;176:106284. Epub 20201210. doi: 10.1016/j.rmed.2020.106284. PubMed PMID: 33338874.
28. Bloch F, Thibaud M, Dugue B, Breque C, Rigaud AS, Kemoun G. Episodes of falling among elderly people: a systematic review and meta-analysis of social and demographic predisposing characteristics. *Clinics (Sao Paulo)*. 2010;65(9):895-903. doi: 10.1590/s1807-59322010000900013. PubMed PMID: 21049218; PubMed Central PMCID: PMC2954741.
29. Sousa LM, Marques-Vieira CM, Caldevilla MN, Henriques CM, Severino SS, Caldeira SM. Risk for falls among community-dwelling older people: systematic literature review. *Rev Gaucha Enferm*. 2017;37(4):e55030. Epub 20170223. doi: 10.1590/1983-1447.2016.04.55030. PubMed PMID: 28273251.
30. Singh T, Belanger E, Thomas K. Is Fear of Falling the Missing Link to Explain Racial Disparities in Fall Risk? Data from the National Health and Aging Trends Study. *Clin Gerontol*. 2020;43(4):465-70. Epub 2018/05/17. doi: 10.1080/07317115.2018.1468377. PubMed PMID: 29764333; PubMed Central PMCID: PMC6202265.
31. Iezzoni LI, Heeren T, Foley SM, Daley J, Hughes J, Coffman GA. Chronic conditions and risk of in-hospital death. *Health Serv Res*. 1994;29(4):435-60. PubMed PMID: 7928371; PubMed Central PMCID: PMC1070016.

32. Heron M. Deaths: Leading Causes for 2016. *Natl Vital Stat Rep.* 2018;67(6):1-77. PubMed PMID: 30248017.
33. Utter GH, Cox GL, Atolagbe OO, Owens PL, Romano PS. Conversion of the Agency for Healthcare Research and Quality's Quality Indicators from ICD-9-CM to ICD-10-CM/PCS: The Process, Results, and Implications for Users. *Health Serv Res.* 2018;53(5):3704-27. Epub 2018/05/31. doi: 10.1111/1475-6773.12981. PubMed PMID: 29846001; PubMed Central PMCID: PMC6153152.
34. Tinetti ME, McAvay GJ, Fried TR, Allore HG, Salmon JC, Foody JM, et al. Health outcome priorities among competing cardiovascular, fall injury, and medication-related symptom outcomes. *J Am Geriatr Soc.* 2008;56(8):1409-16. Epub 2008/07/30. doi: 10.1111/j.1532-5415.2008.01815.x. PubMed PMID: 18662210; PubMed Central PMCID: PMC3494099.
35. Pisani L, Hill NS, Pacilli AMG, Polastri M, Nava S. Management of Dyspnea in the Terminally Ill. *Chest.* 2018;154(4):925-34. Epub 2018/04/22. doi: 10.1016/j.chest.2018.04.003. PubMed PMID: 29679597.
36. Currow DC, Agar MR. Benzodiazepine Prescribing in People with Chronic Obstructive Pulmonary Disease: Clinical Considerations. *Drugs Aging.* 2020;37(4):263-70. Epub 2020/02/29. doi: 10.1007/s40266-020-00756-z. PubMed PMID: 32107742.
37. Racey M, Markle-Reid M, Fitzpatrick-Lewis D, Ali MU, Gagne H, Hunter S, et al. Fall prevention in community-dwelling adults with mild to moderate cognitive impairment: a systematic review and meta-analysis. *BMC Geriatr.* 2021;21(1):689. Epub 2021/12/10. doi: 10.1186/s12877-021-02641-9. PubMed PMID: 34893027; PubMed Central PMCID: PMC8665555.
38. Peek K, Bryant J, Carey M, Dodd N, Freund M, Lawson S, et al. Reducing falls among people living with dementia: A systematic review. *Dementia (London).* 2020;19(5):1621-40. Epub 2018/09/30. doi: 10.1177/1471301218803201. PubMed PMID: 30269532.
39. McDermott CL, Engelberg RA, Woo C, Li L, Fedorenko C, Ramsey SD, et al. Novel Data Linkages to Characterize Palliative and End-Of-Life Care: Challenges and Considerations. *J Pain Symptom Manage.* 2019. doi: 10.1016/j.jpainsymman.2019.07.017. PubMed PMID: 31349037.

Table 1. Demographics of 8,204 patients with COPD by injurious fall status

	Had injurious fall N=2454	No injurious fall N=5750
Mean age (SD)	70.1 (13.3)	70.2 (12.7)
Male (n, %)	1498 (61)	3523 (61)
Female (n, %)	956 (39)	2227 (39)
Race (n, %)		
White, non-Hispanic	1916 (78)	4838 (84)
Asian	113 (5)	248 (4)
Black	272 (11)	345 (6)
White, Hispanic	51 (2)	104 (2)
Native American	60 (2)	98 (2)
Pacific Islander	8 (0.5)	26 (0.5)
Multiracial	32 (1)	71 (1)
Unknown	2 (0.5)	20 (0.5)
Married (n, %)	604 (25)	2277 (40)
Some college (n, %)	954 (39)	2466 (43)
Insurance (n, %)		
Private	279 (11)	995 (17)
Medicare	1194 (49)	3127 (54)
Medicaid	746 (30)	1078 (19)
Military/Other	126 (5)	298 (5)
Unknown	109 (5)	252 (5)

Table 2, Comorbidities and fall history among 8,204 patients with COPD, by injurious fall ^a

N (%)	Had injurious fall N=2454	No injurious fall N=5750
Fall history	576 (23)	244 (4)
Cancer	645 (26)	2039 (35)
Coronary artery disease	1213 (49)	2625 (46)
Dementia	485 (20)	511 (9)
Diabetes	490 (20)	869 (15)
End-stage liver disease	370 (15)	505 (9)
Heart failure	1152 (47)	2352 (41)
Other pulmonary disease	1038 (42)	2257 (39)
Peripheral vascular disease	731 (30)	1544 (27)
Renal disease	783 (32)	1457 (25)
Total comorbidities (mean, SD)	2.81 (1.80)	2.46 (1.56)

^a Patients may have more than one comorbidity, so comorbidity totals are greater than the total number of patients.

Table 3, FRID prescription history among 8,204 patients with COPD by injurious fall ^a

N (%)	Had injurious fall N=2454	No injurious fall N=5750	P-value
Anticonvulsants	857 (35)	1497 (26)	<0.05
Antihypertensives	1578 (64)	3740 (65)	0.52
Antipsychotics	597 (24)	768 (13)	<0.05
Antispasmodics	742 (30)	1449 (25)	<0.05
Atypical antidepressants ^b	683 (28)	1102 (19)	<0.05
Benzos	970 (40)	1961 (34)	<0.05
Opioids	1673 (68)	3686 (64)	<0.05
Tricyclic antidepressants	242 (10)	291 (5)	<0.05
Sedative Hypnotics ^c	281 (11)	433 (8)	<0.05
SNRIs	214 (9)	316 (5)	<0.05
SSRIs	671 (27)	1128 (20)	<0.05

^a Patients may have more than one prescription, so prescription totals are greater than the total number of patients.

^b (e.g. bupropion, mirtazapine)

^c (e.g. eszopiclone, zolpidem)

Online Supplement

Pre-proof